

Everyone who is in any way involved with a disaster or traumatic event may experience trauma reactions. You might experience these reactions if you:

- Were a witness or were involved in the event
- Arrived upon the scene of the event
- Had a “near miss” or were almost involved in the event
- Knew or know others who were killed, harmed, or involved in some way
- Have a relationship with family or friends of victims.
- Are reminded of other traumatic incidents in your life by this event.

It is important to remember that trauma reactions are normal reactions to extremely abnormal circumstances. It is difficult to predict what type of trauma reactions you will experience following a disastrous event. It is important to allow yourself permission to have your reactions, and take care of them both by yourself and by asking for help from others, as best you can. Many people find it helpful to have information about what constitutes a typical reaction to trauma.

## TYPICAL REACTIONS TO TRAUMA

Not everyone experiences the same set of responses to trauma, but people typically experience reacts that fall into four basic categories. Here are some reacts that you may be experiencing:

### PSYCHOLOGICAL AND EMOTIONAL

- Heightened anxiety or fear
- Irritability, restlessness, or over excitability
- Feelings of sadness, moodiness, ore crying then usual
- Feeling of helplessness or hopelessness
- Felling of numbness or detachment
- “Survivor guilt”, or feeling of self-blame that you escaped the tragedy
- Re-experiencing of traumatic event, possibly including:
  - Intrusive thought or images of the event
  - Distressing dreams or nightmares
  - Flashbacks about the event
  - Distress when exposed to events that remind you of trauma
- Feelings of estrangement or isolation from others
- Hyper-vigilance (feelings especially attuned to events around you, scanning environment for possible danger)

### COGNITIVE

- Difficulty concentrating
- Feeling confused or distracted, slower thought than normal

### PHYSICAL

- Headaches
- Nausea or upset stomach
- Exaggerated startle response (tendency to startle easily at loud noises)
- Fatigue or feeling slowed down

### BEHAVIORAL

- Hyperactivity, or less activity
- Heightened tendency to behave irritably
- Withdrawal, social isolation
- Avoidance of activities or places that remind you of traumatic event
- Insomnia or sleeping excessively
- Strong need to talk about the event, read accounts about the event

You may recognize yourself as experiencing some of the above reactions. Remember that your response is normal. Immediately following a traumatic event you will probably feel disrupted, dazed, and somewhat confused. You will notice that you are not behaving as you typically would. It is important to take care of yourself as best you can. Here are some self-care suggestions for you.

## SELF CARE

1. Keep reminding yourself that your responses are normal responses to a stressful situation. Give yourself permission to do whatever you need to do to take care of yourself. Your body and mind will tell you what you need to do – your job is to listen to them.
2. Get plenty of rest when you're tired, and use the energy you have if you experience hyperactivity at times. Don't force yourself to be active if you don't have the energy, or rest when you don't feel tired.
3. Talk to people as much as you need to. Reach out. You may experience a need to talk repetitively about the trauma. If you can find someone who is willing to listen, use her/him to talk to about how you are feeling. If you don't have anyone in your support network to use, talk to your medical provider or behavioral health consultant who can refer you to someone.
4. Spend time with others, even if you don't feel like talking. It can be very comfortable to know you're not alone. Try to find someone or someplace that feels safe and comforting to you, and spend time there.
5. Don't make any major life decision or big life changes if at all possible. This is not a time to put pressure on yourself to do anything out of the ordinary. Concentrate on taking care of yourself.
6. Do things that feel good to you – take baths, read, exercise, watch television, spend time with friends and family, fix yourself a special treat, or whatever else feels nurturing and self caring.
7. Allow yourself to cry, rage, and express your feeling when you need to. Try not to numb your feelings with alcohol or drugs. This will only complicate your situation.

## THE PROCESS OF RECOVERY

It is important to know that recovering from a trauma is a process that may take a long time. The initial response of disruption (perhaps alternating with numbness) may last days, weeks, or longer. Don't be surprised if you continue to experience these reactions for longer than you expected. It is impossible to predict how long you will experience effects of the trauma, but usually trauma reactions gradually decrease over time. If you experience another stressful event while recovering from trauma, you may find that your trauma reactions reappear for a while. This re-activation, or delayed trauma response, is perfectly normal.

At any time during this process, you may find it useful to ask for professional help from a counselor or mental health professional. There are some circumstances under which you should definitely get professional help:

If you find yourself feeling suicidal or contemplating suicide

If you find that your daily function continues to be impaired so that you cannot carry out your life tasks.

If your post-trauma fears interfere with your ability to return to certain places or situations that remind you of the trauma.

## IMPORTANT PHONE NUMBERS

Marin Community Clinics: 415-448-1500

24-hour Suicide Prevention Line: 415-499-1100

Marin County Resources: 2-1-1

Marin County Behavioral Health Services: 1-888-818-1115

Mobile Crisis Unit: 415-473-6392 (1pm-9pm)