Patient's Name:	Date of Birth:		
Worksheet for DSM-V criteria for diagnosis of	opiat	e use d	isorder
Diagnostic Criteria* (Opioid Use Disorder requires at least 2 within 12 month period)	Meets criteria Yes No		Notes/supporting information
Opioids are often taken in larger amounts or over a longer period of time than intended.			
There is a persistent desire or unsuccessful efforts to cut down or control opioid use.			
3. A great deal of time is spent in activities necessary to obtain the opioid, use the opioid, or recover from its effects.			
4. Craving,or a strong desire to use opioids.			
<ol> <li>Recurrent opioid use resulting in failure to fulfill major role obligations at work, school or home.</li> </ol>			
<ol><li>Continued opioid use despite having persistent or recurrent social or interpersonal problems caused or exacerbated by the effects of opioids.</li></ol>			
7. Important social, occupational or recreational activities are given up or reduced because of opioid use.			
Recurrent opioid use in situations in which it is physically hazardous			
<ol> <li>Continued use despite knowledge of having a persistent or recurrent physical or psychological problem that is likely to have been caused or exacerbated by opioids.</li> </ol>			
<ul><li>10. *Tolerance, as defined by either of the following:</li><li>(a) a need for markedly increased amounts of opioids to achieve intoxication or desired effect</li><li>(b) markedly diminished effect with continued use of the same amount of an opioid</li></ul>			
<ul><li>11. *Withdrawal, as manifested by either of the following:</li><li>(a) the characteristic opioid withdrawal syndrome</li><li>(b) the same (or a closely related) substance are taken to relieve or avoid withdrawal symptoms</li></ul>			
*This criterion is not considered to be met for those individuals	taking	opioids	solely under

\*I his criterion is not considered to be met for those individuals taking opioids solely under appropriate medical supervision.

Severity: Mild: 2-3 symptoms, Moderat	e: 4-5 symptoms. <b>Severe</b> : 6 or more symptoms.
Signed	Date

Criteria from American Psychiatric Association (2013). <u>Diagnostic and Statistical Manual of Mental Disorders</u>, Fifth Edition, Washington, DC, American Psychiatric Association page 541.